



URINARY SUPPORT:

GRAMINEX Flower Pollen Extract

Inhibitory Effect and Synergism of Cernitin™ Pollen Extract on the Urethral Smooth Muscle and Diaphragm of the rat

Nakase K, Takenaga K, Hamanaka T, Kimura M

Inhibitory effects of Cernitin™ pollen extract (CN-009), consisting of T-60 (a water-soluble extract) and GBX (an acetone-soluble extract) at a ratio of 20:1, were investigated in rat urethral smooth muscle and diaphragm. In the urethral smooth muscle, CN-009 (3.0×10^{-4} approximately 3.0×10^{-3} g/ml), T-60 and GBX (corresponding to CN-009) concentration-dependently inhibited the noradrenaline (NA, 10^{-5} g/ml)-induced contraction. According to Burgi's calculation, the inhibition by T-60 and GBX was synergistic. On the other hand, GBX inhibited the NA-contraction non-competitively and also inhibited the K⁺-contraction. In contrast, T-60 tended to inhibit competitively, but did not affect the K⁺-contraction. In the diaphragm, CN-009 (5.25×10^{-3} approximately 2.1×10^{-2} g/ml) concentration-dependently inhibited the nerve stimulation-induced twitch response. T-60 (corresponding to CN-009) showed no effect, while GBX slightly inhibited the twitch response. The effects of T-60 and GBX were synergistic. The inhibitory effects of CN-009 (2.1×10^{-2} g/ml) and GBX (1.0×10^{-2} g/ml) were specific against the nerve stimulation and were not antagonized by neostigmine (1.0×10^{-5} g/ml). These results suggested that these effects were neither musculotropic nor competitive against ACh. In conclusion, CN-009 concentration-dependently inhibited the urethral contraction and the skeletal muscular twitch response. It was suggested that T-60 and GBX had different mechanisms and inhibited the response synergistically.

PMID: 3417212, UI: 88329868

Ome Research Laboratories Tobishi Pharmaceutical Co., Ltd.

Tokyo, Japan.

Nippon Yakurigaku Zasshi 1988 Jun;91(6):385-92