



# Demonstration of superoxide dismutase enzymes in extracts of pollen and anther of *Zea mays* and in two related products, Baxtin® and Polbax®

PER CHRISTER ODÉN, GÖSTA KARLSSON and ROLAND EINARSSON

Odén, P.C., Karlsson, G. & Einarsson, R. 1992. Demonstration of superoxide dismutase enzymes in extracts of pollen and anther of *Zea mays* and in two related products, Baxtin® and Polbax®. – Grana 31: 76-80, 1992. Odense, February 1992. ISSN 0017-3134.

Superoxid dismutase enzymes were isolated and identified in an aqueous extract of pollen and anthers of *Zea mays* and in two commercial products, Baxtin® and Polbax®, derived from the same kind of source material. Prior to analysis the samples were purified by adding soluble poly-N-vinylpyrrolidon and precipitated with ammonium sulphate.

Gel filtration chromatography using a Superose 12 HR column gave well-resolved and similar elution patterns for the maize extract, Baxtin and Polbax samples. Fractions exhibiting superoxide dismutase activity, determined by direct  $KO_2$  assay, were combined, dialysed and evaporated to dryness. The molecular weight of these fractions was approximately 30000 d. These fractions were also analyzed by native polyacrylamide gel electrophoresis and stained for superoxide dismutase enzyme activity using nitro blue tetrazolium. The major region of superoxide dismutase enzyme activity was inhibited by addition of cyanide and hydrogen peroxide indicating the presence of a copper, zinc superoxide dismutase. Another minor region of enzyme activity, migrating as standard manganese superoxide dismutase and not inhibited by cyanide or hydrogen peroxide, was also detected.

The results clearly demonstrated the occurrence of both copper, zinc- and manganese superoxide dismutase enzymes in extracts of pollen and anthers of *Zea mays* and also in the two related commercial products, Baxtin and Polbax.

*Per Christer Odén, Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, S-901 83 Umeå; Gösta Karlsson, PL 4050, S-262 94 Ångeholm; Roland Einarsson, Pharmacia Diagnostics AB, S-751 82 Uppsala, Sweden.*

*(Manuscript received 1 May 1991; revised version accepted 6 August 1991)*

Active oxygen in biological systems has been thoroughly studied and frequently reviewed (Elstner 1987, Salin 1987, Alscher & Amthor 1988, Leshem 1988, Winston 1990). The harmful effect of active oxygen can be reduced by keeping the  $O_2$ -concentration low within the cell through oxidative metabolism or oxygnease reactions. Other possibilities are to keep

substrates that produce toxic interme diates separated from each other or by scavenging systems, antioxidants, that reduce the concentrations of active oxygen species. The antioxidants can either be fat-soluble, e.g., carotenoids and  $\alpha$ -tocopherol, water-soluble, e.g., glutathione and ascorbic acid, or enzymatic

compounds, e.g., glutathione reductase and superoxide dismutase (SOD).

SOD is a metalloenzyme or actually a group of metalloenzymes that catalyzes the disproportionation of  $O_2^-$  to yield molecular  $O_2$  and  $H_2O_2$  (Weselake et al. 1986). SOD protects cells against oxygen toxicity by scavenging the superoxide radical and SOD is therefore a key component in the free radical detoxification process. SOD-enzymes have been extensively studied during the last ten years and are now considered present in all aerobic organisms (Rabinowitch & Fridovich 1983, Fridovich 1986, Asada 1988, Monk et al. 1989, Hassan & Scandalios 1990). Three types of SODs have been found in plants, classified according to the metal at the catalytic centre: copper and zinc SOD (CuZn SOD), a manganese-containing homo dimer (Mn SOD) and an iron-containing homo dimer (Fe SOD) (Fridovich 1986). These three types can be distinguished by their sensitivity to inhibitors, CuZn SOD is inhibited by cyanide and  $H_2O_2$  and the Fe SOD is inhibited by  $H_2O_2$ , whereas the Mn SOD is insensitive to both inhibitors. Most CuZn SODs isolated so far, with few exceptions, are homodimers with a molecular weight of approximately 32,000 (Fridovich 1986). The most abundant SODs in

plants are the CuZn SODs, which are found mainly in the cytosol and chloroplasts. Mn SOD is usually localized to the mitochondrial matrix in higher plants (Asada 1988) but has also been isolated from chloroplast thylakoids (Hayakawa et al. 1985). In addition it has been detected in the glyoxysomes (Del Rio et al. 1983, Sandalio & Del Rio 1987).

The development of reproductive structures, e.g., pollen and spores, and the pollination and critical phases in the life cycle of plants. As good protection against the detrimental effects of free radicals is therefore necessary to avoid damage to the genetic information. Recently, Acvedo & Scandalios (1990) reported on the expression of SOD genes in mature pollen of maize.

The purpose of the present investigation was to demonstrate the presence of SOD enzymes in three different extracts of pollen and anthers of maize (*Zea mays*); freshly prepared maize extract, Baxtin and Polbax.




Fig. 1. Superose 12 gel filtration pattern of the maize pollen and anthers extract. The marked area contained SOD activity. Arrows indicate retention volumes of the molecular weight markers and the standard SOD enzymes; thyroglobulin (1), ferritin (2), catalase (3), aldolase (4), ovalbumin (5), chymo-trypsinogen A (6), CuZn SOD (7), Mn SOD (8) and Fe SOD (9).

## MATERIALS AND METHODS

### *Preparation and purification of maize extract, Baxtin-solution and dissolved Polbax tablets*

Fresh pollen and anthers of *Zea mays* (ratio 10:1 w/w) were incubated for 18 h in 200 ml of 50 mM TRIS-HCl, pH 7.5, containing 5 mM MgCl<sub>2</sub>, 10 mM 2-mercapto-ethanol, 0.4 mM ascorbate, 2 mM EDTA and 4% poly-N-Vinylpyrrolidone 360 (PVP) (w/w). After incubation the sample was homogenized in an ice-bath with an Ultra-Turrax at maximum speed for 5 min. The tissue debris was removed by filtering and the sample was carefully brought to 20% saturation with saturated ammonium sulphate solution, stirred for 30 min and the precipitate was removed by centrifugation at 5000 X g for 30 min. The supernatant was then brought to 90% saturation with addition of solid ammonium sulphate and stirred for 1 h, The precipitate was collected by centrifugation at 5000 X g for 30 min, and dissolved in 2.5 ml of 10 mM TRIS-HCl, pH 7.0. The SOD activity in this extract was tested by the direct KO<sub>2</sub> assay as described below and thereafter isolated using high-performance liquid chromatography equipped with a Superose TM 12 HR 10/30 column (highly cross-linked agarose matrix, Pharmacia LKB Biotechnology, Uppsala, Sweden). The sample was injected off-column via an injector with a 200 µl loop and the column was eluted with 5 mM phosphate buffer containing 0,15 m NaCl at a flow rate of 0.75 ml per min. The absorbance of the eluate was monitored at 254 nm. Forty fractions of 0.75 ml each were collected and the fractions were tested for SOD activity using the direct KO<sub>2</sub> assay.

For calibration of the Superose gel filtration column the elution volumes of the following molecular weight markers was determined; thyroglobulin 669.000 d, ferritin 440.000 d. catalase 232.000 d, aldolase 158.000 d, ovalbumin 43.000 d, chymotrypsinogen A 25.000 d (Pharmacia LKB Biotechnology) and standard SOD enzymes; CuZn SOD from

Horseradish, Mn SOD from *E. coli* and Fe SOD from *E. coli* (Sigma Chemical Company, St. Louis, MO, USA).

Baxtin is a commercially available extract solution of fresh and well-controlled pollen and anthers of *Zea mays* (ratio 10:1 w/w) manufactured by Allergon AB, Välinge, Sweden. Ten ml of Baxtin solution (batch no 97808101) was carefully added in drops while stirring to 190 ml of 50 mM TRIS-HCl, pH 7.5, containing 5 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, 0.4 mM ascorbate, 2 mM EDTA and 4% PVP 360 (w/w). The Baxtin solution was then treated as described above for the freshly prepared maize extract of pollen and anthers.

Polbax (Allergon AB) is a commercially available product produced from a specified amount of grass pollen extract (Pollixin) and Baxtin. Polbax (210 tablets, batch no Q 186218) were ground to a fine powder and dissolved in 200 ml of 50 mM TRIS-HCl, pH 7.5, containing 5 mM EDTA and 4% PVP 360 (w/w). The Polbax solution was thereafter treated in an analogous manner to the previously described maize extract.

### *Determination of SOD activity by the direct KO<sub>2</sub> assay*

The SOD enzyme activity in the sample was measured by analyzing the disproportionation of O<sub>2</sub><sup>•-</sup>. according to Marklund (1982). The analyzed samples were added in volumes of 20-40 µl to a 1 cm quartz cuvette containing 3 ml of 50 mM 2-amino-2-methyl-1 propanol (AMP)-HCl plus 0.2 mM diethylenetriamine penta acetic acid (DTPA), pH 9.5 and 5 µl 30 µM bovine liver catalase. KO<sub>2</sub> was dissolved in ice cold 50 mM NaOH plus 0.5 mM DTPA and 15 µl of this solution was added as substrate to the reaction mixture. The decay of the added O<sub>2</sub><sup>•-</sup> was then measured continuously until the baseline had stabilized. KO<sub>2</sub> was added several times to confirm the enzymatic nature of the disproportionation reaction of the analyzed samples. The sensitivity of the disproportionation of O<sub>2</sub><sup>•-</sup> to CN<sup>-</sup> was also

monitored by adding 30  $\mu$ l of 0.3 M NaCN to the reaction mixture.

### *Polyacrylamide gel electrophoresis (PAGE) and NBT staining for SOD activity*

Native PAGE was performed at 15° C using PhastSystem and PhastGel homogenous 20 (Pharmacia LKB Biotechnology). Pooled fractions containing SOD activity from the gel filtration chromatography were dialysed for 24 h against 4 l of 5 mM TRIS-HCl buffer pH 7.0 and finally evaporated to dryness. The samples were thereafter dissolved in 10 mM TRIS-HCl buffer, pH 7.0 and applied on native PAGE. The standard SOD enzymes were run in parallel.

The gels were immediately stained for SOD enzyme activity according to Beauchamp & Fridovich (1971) after finished electrophoresis. The gels were first soaked in darkness in 2.45 mM nitro blue tetrazolium (NBT) for 20 min and then soaked in 28 mM tetramethyl ethylene diamine plus 28  $\mu$ M riboflavin for 15 min. All reagents were dissolved in 36 mM phosphate buffer pH 7.8. The effect of inhibitors in the activity was studied by adding 10 mM NaCN and 10 mM H<sub>2</sub>O<sub>2</sub>, respectively, to the solutions. After staining in darkness the gels were carefully washed with water and then treated with light. Regions with SOD enzyme activity were then visualized as lighter parts on the darkblue gel.




Fig. 2. Disproportionation of O<sub>2</sub><sup>-</sup> by a fraction of the pollen and anthers extract measured spectrophotometrically at 250 nm after various additions of KO<sub>2</sub> and the effect of CN<sup>-</sup> on the disproportionation of O<sub>2</sub><sup>-</sup>.

## RESULTS AND DISCUSSION

SOD enzyme activity was measured by the direct KO<sub>2</sub> assay in the crude extract of pollen and anthers of *Zea mays*, Baxtin and dissolved Polbax tablets. Due to analytical disturbances in the samples, however, it was not possible to obtain well-resolved protein bands with defined SOD enzyme activity on native PAGE gels after NBT staining.

Therefore the samples had to be purified to localize the active SOD enzymes in the electrophoretic patterns. The first step in the purification sequence was to absorb interfering compounds, mainly phenols, to soluble PVP. The second step was to apply the maize extract on a Superose 12 HR 10/30 column for further purification by gel filtration chromatography. The gel filtration elution pattern of the pollen and anther extract is shown in Fig. 1. SOD enzyme activity was detected by the direct KO<sub>2</sub> assay in fractions eluting between 13.5 and 15 ml. These active fractions were pooled for further analysis. Fig. 2. shows the O<sub>2</sub><sup>-</sup> disproportionation activity of the fractions eluted at 14.25-15.00 ml versus time after several additions of KO<sub>2</sub>. The repeatability of the activity pattern after KO<sub>2</sub> addition for this maize fraction indicates that




Fig. 3. Superose 12 gel filtration pattern of the Baxtin sample. Experimental conditions as in Fig. 1.

the molecule(s) responsible for the disproportionation reaction are not consumed and thus constitute an enzyme molecule. Furthermore the effect of  $CN^-$ , a Cu ZnSOD enzyme inhibitor, on the  $O_2^-$ -disproportionating activity of the maize fraction is also shown in Fig. 2. The maize-derived pollen and anther fraction exhibits SOD enzyme activity and the main part of this activity is inhibited by the addition of  $CN^-$ . The cyanide sensitivity of the maize fraction confirms that the enzyme is copper and zinc-containing SOD (Fridovich 1986).

Fig. 1 also presents data about elution volumes of the molecular weight markers. The identified SOD enzymes at 2500 d and 43000 d and further the standard SOD enzymes chromatographed in parallel with the maize extract also elutes in this molecular weight region. A linear regression of the logarithmic molecular weights versus retention volumes resulted in a coefficient of determination of 0.97 and an approximate mean molecular weight of the putative SOD enzyme of 28.600.

The gel filtration elution profiles of the purified Baxtin and Polbax samples are shown in Figs. 3 and 4. The Superose elution patterns of Baxtin (Fig. 3) and the freshly prepared maize extract (Fig. 1) are similar, while the pattern for Polbax differs in the low molecular weight region (Fig. 4), most likely due to the presence of grass pollen components (e.g., Pollixin). However the elution volume of the fractions containing SOD enzyme activity is located in the same region. Both Baxtin and Polbax samples exhibited maximum SOD enzyme activity in the fractions eluting between 13.5 ml and 15 ml. The main part of the SOD enzyme activity was also inhibited by the addition of 10 mM  $CN^-$  to the reaction medium when performing the direct  $KO_2$  assay, confirming the presence of CuZnSOD in Baxtin and Polbax.

The homogeneity of the SOD enzyme activity in the pooled fraction from Superose gel filtration was analyzed by PAGE. The polyacrylamide gel




Fig. 4. Superose 12 gel filtration pattern of the Polbax sample. Experimental conditions as in Fig. 1.

used allow complex mixtures of proteins to be separated in the molecular weight range 2000-150000 d. Fig. 5 illustrates three native PAGE gel patterns with standard CuZn SOD, FeSOD, MnSOD and the pooled fractions (13.5-15.0 ml) of the freshly prepared maize extract, Baxtin and Polbax samples in the absence and presence of SOD enzyme inhibitors during the staining procedure. The SOD enzyme activity in the various maize samples applied on the gel was developed by using the NBT reagent.

Fig. 5A demonstrates distinct SOD enzyme activity for all three analyzed maize samples (lane b, c, d) and further the activity is localised to defined regions on the gel. The enzyme activity of the SOD standards run in parallel is easily localised in the PAGE gel pattern. The enzyme activity of standard CuZn SOD and MnSOD partly overlap the SOD enzyme activity of the maize samples. Fig. 5B shows a PAGE analysis where the gel was treated with 10 mM  $CN^-$  and Fig. 5C with 10 mM  $H_2O_2$  during the staining procedure. Inactivation of the enzyme activity (lane b, c, d) in the presence of  $CN^-$  confirmed that the SOD activity was due to CuZnSOD (Fig. 5B). The region located closer to the front was clearly inhibited by the addition of  $CN^-$  and  $H_2O_2$  as was also the standard CuZn SOD (Fig. 5B and 5C). On the contrary the region close to the application point on the




Fig. 5. A-C. Native PAGE gels stained for SOD enzyme activity with NBT after analysis of standard CuZn SOD (500 ng) (a), pollen and anthers extract (b), Baxtin sample (c), Polbax sample (d), standard Mn SOD (250 ng) (e) and standard Fe SOD (250 ng) (f). Polyacrylamide gels; (A) without inhibitor (B) in the presence of 10 nM  $CN^-$  during staining and (C) and in the presence of 10 mM  $H_2O_2$  during staining.

PAGE gel was neither inhibited by  $CN^-$  nor  $H_2O_2$  and migrated as standard Mn SOD (lane e), therefore indicating the presence of a Mn SOD in the samples. The activity of the standard Fe SOD (lane f) was inhibited by the addition of  $H_2O_2$  but not by addition of  $CN^-$  (Fig. 5B and 5C). No further change in the SOD enzyme activity pattern of maize pollen and anther extract, Baxtin or Polbax samples could be detected after  $H_2O_2$  treatment (Fig. 5C) demonstrating that FeSOD is not present in the analyzed maize extracts. The NBT activity staining patterns are in agreement with the gelfiltration data i.e. the detected SOD enzyme activity in the analyzed maize preparations are located in the regions for low molecular weight protein molecules.

## CONCLUSIONS

The presence of superoxide dismutase enzymes in extracts of pollen and anthers of *Zea mays* and in the related products Baxtin and Polbax was established after purification of the samples by a simple PVP precipitation. Superose gel filtration of the purified samples revealed that SOD enzyme activity was located around 30000 d. The enzyme activity was inhibited by  $CN^-$  indicating the presence of CuZnSOD. Polyacrylamide gel electrophoresis followed by NBT activity staining in the absence and presence of different inhibitors confirmed that the SOD enzyme activity in the pooled fractions from Superose gelfiltration of the freshly prepared maize extract, Baxtin and Polbax samples was due to CuZnSOD and MnSOD.

## REFERENCES

- Acvedo, A. & Scandalios, J. G. 1990. Expression of the catalase and superoxide dismutase genes in mature pollen in maize. – *Theor. Appl. Genet.* 80: 705-711.
- Alscher, R. G. & Anthor, J. S. 1988. The physiology of free radical scavenging: Maintenance and repair processes. – In: *Air pollution and plant metabolism* (ed. S. Shulte-Hostede, N. M. Darral, L. W. Blank & A.R. Wellburn). pp. 94-115. – Elsevier, London – New York.
- Asada, K. 1988. Superoxide dismutase. – In: *Metalloproteins* (ed. S. Otsuka & T. Yamanaka). pp. 331-341. – Elsevier, Amsterdam.
- Beauchamp, C. & Fridovich, I. 1971. Superoxide dismutase: Improved assay and an assay applicable to acrylamide gels. – *Anal. Biochem.* 44: 276-287.
- Del Rio, L. A., D. S., Olah, I., Glick, B. & Salin, M. L. 1983. Immunocytochemical evidence for a peroxisomal localization of manganese superoxide dismutase in leaf protoplasts from a higher plant. – *Planta* 158: 216-224.
- Elstner, E. F. 1987. metabolism of activated oxygen species. – In: *chemistry of metabolism*. Vol. 11 (ed. P.K. Strumpf & E. E. Conn). pp. 253-315. – Acad. Press, New York.
- Fridovich, I. 1986. Superoxide dismutases. – In: *Advances in Enzymology*. Vol. 58 (ed. A. Meister), pp. 61-97. – New York.
- Hassan, H.M. & Scandalios, J. G. 1990. Superoxide dismutase in aerobic organisms. – In: *Stress Responses in plants: Adaptation and acclimation mechanisms* (ed. R. Alscher & J. Cumming), pp. 175-199. – Willey-Liss Inc., New York.
- Hayakawa, T., Kanematsu, S. & Asada, K. 1985. Purification and characterization of thylakoid-bound Mn-superoxide dismutase in spinach chloroplasts. – *Planta* 166: 111-116.
- Leshem, Y. 1988. Plant senescence processes and free radicals. – *Free Radic. Biol. Med.* 5: 39-49.
- Marklund, S. L. 1982. Human copper-containing superoxide dismutase of high molecular weight. – *Proc. Natl. Acad. Sci.* 79: 7634-7638.
- Monk, L. S., Faferstedt, K. V., Crawford, R. M. M. 1989. Oxygen toxicity and superoxide dismutase as an antioxidant in physiological stress. – *Physiol. Plant* 76: 456-459.
- Rabinowitch, H. D. & Fridovich, I. 1983. Superoxide radicals, superoxide dismutases and oxygen toxicity in plants. – *Photochem. Photobiol.* 37: 679-690.
- Salin, M. L. 1987. Toxic oxygen species and protective systems of the chloroplast. – *Physiol. Plant.* 72: 681-689.
- Sandalio, L. M. & Del Rio, L.A. 1987. Localization of superoxide dismutase in glyoxysomes from *Citrullus vulgaris*. Functional implications in cellular metabolism. – *J. Plant. Physiol.* 127: 395-409.
- Weselake, R. J., Chesney, S. L., Petkau, A. & Friesen, A. D. 1986. Purification of human copper, zinc superoxide

dismutase by copper chelate affinity chromatography. –  
Anl. Biochem. 155: 193-197.

17. Winston, G. W. 1990. Physicochemical basis for free radical formation in cells: production and defenses. – In:

Stress responses in plants: Adaptation and acclimation mechanisms (ed. R. Alscher & J. Cumming), pp. 57-86. –  
Wiley-Liss Inc., New York.