



Effect of Cernitin Pollen-Extract on Experimental Nonbacterial Prostatitis in Rats

Toshiyuki Kamijo,¹ Shigeru Sato,² and Tadaichi Kitamura¹

¹Department of Urology, Faculty of Medicine, University of Tokyo, Japan

²Ohme Research Laboratories, Tobishi Pharmaceutical Co., Ltd., Tokyo, Japan

BACKGROUND. The treatment for chronic nonbacterial prostatitis (NBP) has not been established. Cernitin pollen-extract (CN-009) is reported to have therapeutic effects for NBP. The effects and mechanisms of CN-009 were investigated.

METHODS. Ten-month-old rats were used with administration of estradiol after castration, which were similar to human NBP histologically. Since CN-009 consists of T-60 and GBX, these drugs were administered, respectively. The prostate was evaluated histopathologically including glandular damage (epithelial score), stromal ratio and immunohistochemical assays for epithelial function (PAP), stromal evaluation (Vimentin), cell proliferation (PCNA) and apoptosis (deoxyuridine triphosphate biotin nick end-labeling (TUNEL)).

RESULTS. Controls revealed severe acinar gland atrophy and stromal proliferation. CN-009 showed diminished these damages. Epithelial score was better ($P<0.01$) and PAP positive materials were more abundant in CN-009 and GBX than in Controls. The stromal ratio was lower in CN-009 ($P<0.01$) and T-60 ($P<0.05$). There was no difference for PCNA positive cells in the epithelium and stroma, and TUNEL positive cells in epithelium. While, the number of TUNEL positive cells in the stroma of CN-009 and T-60 increase ($P<0.01$).

CONCLUSIONS. These findings suggest that CN-009 protects acinar epithelial cells mainly by GBX and also inhibits stromal proliferation in association with enhanced apoptosis mainly by T-60. *Prostate* 49: 122-131, 2001. © 2001 Wiley-Liss, Inc.

KEYWORDS: cernitin pollen-extract; apoptosis; chronic prostatitis; sex-hormone induced prostatitis

INTRODUCTION

The chronic prostatitis syndromes have been recognized; chronic bacterial prostatitis (CBP), chronic nonbacterial prostatitis (NBP) and prostatodynia. NBP is the most frequent disorder of 64% in these three diseases [1]. The etiology of NBP is unknown. A number of organisms or other factors have been reported to be the possible causes for NBP. They are *Trichomonas vaginalis*, *Chlamydia trachomatis*, genital mycoplasmas, staphylococci, coryneforms, genital viruses [2], biofilms [3], stagnation of prostatic secretion, autoimmune disease, allergy, disorder of sex hormone and psychological effects [4,5]. For the treatment of CBP or NBP, antibiotics of new-quinolone or tetracycline have been administered. However, many cases resist these treatments [6].

CN-009 is a pollen extract, which contains 20:1 ratio of powdered aqueous and organic extract. It is essentially a microbial digest of a standardized mixture of eight plant species grown at the Scania area in southern Sweden. The active ingredients consist of water-soluble (T-60) and fat-soluble (GBX) fractions [7,8]. It was reported that CN-009 showed urine discharge action [9,10], anti-prostatic hypertrophic action [7] and anti-inflammatory effects to the prostate [11] in a preliminary study. Since Ask-Upmark [12] reported CN-009 showed an efficacy to prostatitis, it has been used for the treatment of chronic prostatitis with

high therapeutic effects. However, the mechanisms for these effects remain unknown.

To assess the mechanisms of the anti-prostatitis effect by CN-009, the present study was performed using a nonbacterial prostatitis rat model [13,14] induced by 17 β -estradiol administration and castration.

MATERIALS AND METHODS

Sex Hormone-Induced Nonbacterial Prostatitis Model

Ten month-old Wistar aged male rats were purchased from Japan Slc Co. (Tokyo, Japan). The rats were housed in a climatized environment with a 12-hr light/ dark cycle, 40-70% humidity. Food and water were supplied ad libitum. The rats were castrated under ether anesthesia, and then 17 β -estradiol (Sigma, MI) 0.25 mg/ 2 ml/ kg diluted by sesame oil, as an inducer for prostatitis, was subcutaneously injected into the back of rats for 30 days from 1 day after castration [13,14].

Experimental Structure and Schedule

CN-009 was suspended for 630 or 1,260 mg/ 5 ml with 1% HCO-60 (Japan Surfactant, Tokyo, Japan). T-60 and GBX were similarly prepared for 1,200 and 60 mg/ 5 ml, respectively. Testosterone (TS) (Wako Chemicals, Tokyo, Japan), as a positive control, was diluted for 2.5 mg/ 2 ml with corn oil (Yuro Chemical, Tokyo, Japan).

The experimental structure is shown in Table I and the experimental schedule is illustrated in Figure 1. The rats were divided into seven groups consisting of Sham-operation (Sham-ope), Control, CN-009 630, CN-009 1260, T-60, GBX and TS with five or six animals in each group.

In the Sham-ope group, the rats were treated with only Sham-castration and without any drugs. In the Control group, the rats were injected subcutaneously with 17 β -estradiol for 30 days from the day following castration and administered orally with only 1% HCO-60 5 ml/ kg for 14 days from Day 17. In the CN-009 630, CN-009 1260, T-60 and GBX groups, similar protocols were performed with oral administration of CN-009 630, CN-009 1260, T-60 1200 and GBX 60 mg/ kg, respectively. Also in the TS group, the rats were injected subcutaneously with 17 β -estradiol for 30 days from the next day of castration. After 14 days, TS 2.5 mg/ kg was injected subcutaneously for 14 days. All studies were conducted in accordance with institutional guidelines of animal care and in accordance with Japanese Government Animal Protection and Management Law.

Prostate Weights and Histopathological Evaluation

The rats were sacrificed on the day following the final administration. The prostate was extirpated and weighed. Relative prostatic weight was calculated from body weight and absolute weight.

After fixation in 10% neutral buffered formalin, each prostate was cut into coronal blocks. The tissue samples were dehydrated and embedded in paraffin. Sections (3-4 μ m thickness) were stained with Hematoxyline-Eosin (HE), Periodic acid Schiff (PAS) and Masson's tri-chrome. The specimens were evaluated histopathologically.

Immunohistochemistry

Immunohistochemistry studies were performed with anti-prostatic acid phosphatase (PAP), and Vimentin. PAP staining was performed for the evaluation of glandular epithelial function. In

TABLE I. The Structure of the Experiment

Group	No. of animals	Inflammatory agent	Drug treatment
Sham-ope.	5	No-treatment	No-treatment
Control	6	17 β -estradiol 0.25 mg/kg (s.c.)	1% HCO-60 (p.o.)
CN-009 630	5	17 β -estradiol 0.25 mg/kg (s.c.)	CN-009 630 mg/kg (p.o.)
CN-009 1260	6	17 β -estradiol 0.25 mg/kg (s.c.)	CN-009 1260 mg/kg (p.o.)
T-60	5	17 β -estradiol 0.25 mg/kg (s.c.)	T-60 1200 mg/kg (p.o.)
GBX	6	17 β -estradiol 0.25 mg/kg (s.c.)	GBX 60 mg/kg (p.o.)
TS	5	17 β -estradiol 0.25 mg/kg (s.c.)	Testosterone 2.5 mg/kg (s.c.)

Each parenthesis represents the route of administration. s.c, subcutaneous injection; p.o., oral administration.




Fig. 1. The experimental schedule for this study.

PAP stained specimens, anti-PAP polyclonal antibody (Chemicon International, New York, NY) was diluted by PBS including 0.1% BSA of a 1:100 ratio, and incubated for 2 hr at 37° C. Biotinylated anti-rabbit IgG and the avidin-biotin peroxidase complex (ABC) method was performed. Unitect rabbit immunohistochemistry detection systems (Oncogene Science, New York, NY) were reacted by those methods. Vimentin staining was performed for the evaluation of stromal proliferation. An ImmunoCruz staining system (Santa Cruz BioTech., Santa Cruz, CA) for Vimentin staining was used according to the manufacturer's instructions.

Cell Proliferation and Apoptosis

Cell proliferation and apoptosis were investigated with proliferating cell nuclear antigen (PCNA) and terminal deoxynucleotidyl transferase mediated deoxyuridine triphosphate bitotin nick end-labeling (TUNEL), respectively. PCNA staining was performed with PCNA staining kit (ZYMED Laboratories, South San Francisco, CA). TUNEL method was performed with ApoTag Peroxidase In Situ Apoptosis Detection kit (Intervene, New York, NY). In PCNA and TUNEL specimen, 5,000 cells were counted under a microscope in glandular epithelial cells and stromal cells, respectively.

Acinar Epithelial Score and Stromal Area Ratio

To evaluate glandular damage, acinar epithelial cells were classified and scored, as follows; columnar (2 points), cuboidal (1 point), squamous-like (0 point) shape. Three different pathologists without any information judged the score. Using this scoring evaluation, 20 acinar glands of each specimen were investigated. To assess stromal proliferation, all areas of the specimen and the glandular area were

measured using a digitizer (Graph Tech, Tokyo, Japan) with photomicrographs. Using these findings, the stromal ratio was calculated.

Statistical Analysis

All experiments were repeated at least twice. Each value was demonstrated as the mean±SD. Dunnett's test if in equal variance, or non-parametric Dunnett's test if in unequal variance between treatment groups and Control group was performed after 1-way ANOVA followed by Bartlett variance analysis test. Mann-Whitney U test was performed between the Sham-ope and Control groups.

RESULTS

Body and Prostate Weights (Fig. 2)

There was no significant difference in body weight among the CN-009 630, CN-009 1260, T-60, GBX and TS groups compared with the Control group. Absolute and relative prostate weights were significantly ($P<0.01$) decreased in the Control group compared with the Sham-ope group (Fig. 2) in the CN-009 630, CN-009 1260, T-60 and GBX 60 groups, there was no difference compared with the Control group. In the TS group, absolute and relative prostate weights were very close to the Sham-ope group and were significantly different ($P<0.01$) from other groups.

Histopathology and Immunohistochemistry (PAP and Vimentin Staining)

In the Sham-ope group, the prostate was larger than in other groups. Acinar glands were roundly shaped. The acinar lumen was filled with eosinophilic materials. Acinar epithelial cells were cylindrical with a normally situated nucleus and the supranuclear spaces of these cells




Fig. 2. Relative prostate weight. Each column represents the mean \pm SD. ##Significantly different from the Sham-ope group at $P < 0.01$. **Significantly different from the Control group at $P < 0.01$.

contained secretory materials, which were strongly stained with PAP antibody. A few fibrous tissues were found in the stroma (Figs. 3A and 4A). Vimentin positive cells were few, and the Vimentin positive area was small (data not shown).

In the control group, the prostate was atrophic. Acinar glands were irregularly shaped. The acinar lumen was poor with pale stained eosinophilic materials and filled with inflammatory cell infiltrations mainly characterized by neutrophils. Acinar epithelial cells were flattened similar to a squamous cell. A few secretory materials in the epithelial cells were poorly reacted with PAP antibody. The stroma showed severe proliferation with many lymphocyte and monocyte infiltrations and marked fibrosis with fibroblasts (Figs. 3B and 4B). The stroma was stained very strongly with Vimentin. The Vimentin positive area was significantly increased (data not shown). In the CN-009 630 group, the findings were basically identical with the Control group (data not shown).

In the CN-009 1260 group, acinar glands were more roundly shaped than in the Control group. Acinar epithelial cells were cuboidal, and the supranuclear spaces contained secretory materials stained with anti-PAP that were much more abundant than in the Control group. Inflammatory cell infiltrations into the acinar lumen were diminished. The stroma showed

mild proliferation with a few lymphocytes, monocytes and mild fibrosis with fibroblasts (Figs. 3C and 4C). The Vimentin positive area was much less than that of the Control group (data not shown).

In the T-60 group, acinar epithelial cells were more roundly shaped than in the Control group. Although inflammatory cell infiltrations into the lumen were found, stromal cell infiltrations (Fig. 3D), the Vimentin positive cells were also less than that of the Control group (data not shown).

In the GBX group, acinar epithelial cells were more cuboidal than in the Control group. Epithelial cells contained secretory materials stained with anti-PAP, which was basically identical with the CN-009 1260 group. Diminished cell infiltration into the lumen was found (Fig. 3E). However, the stroma showed a proliferative condition with many lymphocyte and monocyte infiltrations and marked fibrosis with many fibroblasts. The stroma was stained strongly with Vimentin, and the positive area was markedly increased (data not shown).

In the TS group, acinar glands were roundly shaped. The acinar lumen was filled with eosinophilic materials with a few cell infiltrations. Acinar epithelial cells were cylindrical and the supranuclear spaces contained many secretory materials with reactive anti-PAP. However, the stroma was stained strongly with Vimentin and showed mild proliferation with fibroblasts (data not shown).



Fig. 3. HE staining of the prostate in experimental nonbacterial prostatitis rat. (A) Sham-ope group: The acinar lumen is filled with eosinophilic materials without any cells. Acinar epithelial cells are cylindrical. A few fibrous tissues are found in the stroma. (B) Control group: The acinar lumen is filled with induced inflammatory cells mainly characterised by neutrophils. Acinar epithelial cells are flattened similar to squamous cells. The stroma shows severe proliferation with many lymphocyte and monocyte infiltrations and remarkable fibrosis with fibroblasts. (C) CN-009 1260 group: Acinar epithelial cells are cuboidal. Inflammatory cell infiltrations into the acinar lumen are diminished. The stroma shows mild proliferation with a few lymphocytes, monocytes and fibroblasts. (D) T-60 group: Stromal proliferation is relatively mild without severe inflammatory cells. (E) GBX group: Acinar epithelial cells are cuboidal, and diminished inflammatory cell infiltrations are shown. 400x. The bar indicates 100 μ m.

Cell Proliferation and Apoptosis (PCNA and TUNEL Positive Cell Counts (Fig. 5))

No significant differences among the groups were observed in the PCNA positive cell counts in epithelial cells (Fig. 6) or in stromal cells (Fig. 7). In the Sham-ope group, a few TUNEL positive cells were found (Fig. 5A). The findings of the Control group were basically identical with the Sham-ope group (Fig. 5B). In the CN-009 1260 group, TUNEL positive cells in the stroma were more abundant than in the Sham-ope and Control groups (Fig. 5C). In TUNEL positive cell counts, no significant differences were observed in acinar epithelial cells (Fig. 8). However, in the stroma, TUNEL positive cells were significantly ($P < 0.05$) increased in the CN-009 1260 group or

T60 group compared with the Control group (Fig. 9).

Acinar Epithelial Score (Fig. 10)

In the Control group, acinar epithelial score was significantly lower ($P < 0.01$) than that of the Sham-ope group. In comparison with the Control group (Fig. 10), the acinar epithelial score was significantly higher ($P < 0.01$) in the CN-009 1260, GBX, and TS groups.

Stromal Area Ratio (Fig. 11)

In the Control group, the stromal area ratio was significantly higher ($P < 0.01$) than that of Sham-ope group in comparison with the Control group.




Fig. 4. Immunohistochemical findings (PAP staining) of the prostate in experimental nonbacterial prostatitis rats. **(A)** Sham-ope group: Supranuclear spaces of acinar epithelial cells contain secretory materials which are stained with anti-PAP. **(B)** Control group: Acinar epithelial cells are flattened similar to squamous cell. Secretory materials are poorly reactive with anti-PAP. **(C)** CN-009 I260 group: Supranuclear spaces contained secretory materials with PAP staining, which are significantly more abundant than in the Control group. $\times 400$ The bar indicates 10 μm .

In comparison with the Control group (Fig. 11) the stromal area ratio of the CN-009 I260 was significantly ($P < 0.01$) lower. The T-60 group was also significantly ($P < 0.05$) lower than the Control group. However, there was no difference between other groups.

Discussion

Although chronic prostatitis is a common disease, it is very difficult to treat effectively.

Typical clinical findings are decreased potential, perineal or scrotal pain, urethral discharge and lower urinary tract irritative symptoms. The prostate gland is irregularly indurated and the numbers of leukocytes in expressed prostatic secretion are increased [15]. Pathological findings of this disease are chronic inflammation characterized by aggregates of lymphocytes in the stroma and




Fig. 5. Immunohistochemical findings (TUNEL) of the prostate in rats. **(A)** Sham-ope group: A few TUNEL positive cells are shown. **(B)** Control group: The findings are basically identical to these of the Sham-ope group. **(C)** CN-009 I260 group: TUNEL positive cells in the stroma are more abundant compared with the Sham-ope and Control groups. $\times 400$. The bar indicates 10 μm .




Fig. 6. Effects of CN-009 on acinar epithelial PCNA positive cell counts of the prostate. Each column represents the mean \pm SD.

acute inflammation characterized by the presence of neutrophilic polymorphonuclear leukocytes in the lumen of acinar glands [15-17]. Pathological definition of chronic prostatitis is different from clinical definition for the urologists. Clinical definition has been the combination of a clinical symptom and inflammatory cells in expressed prostatic secretion. The pathological inflammation of the prostate was reported to be not frequent in patients with symptoms of chronic prostatitis/ chronic pelvic pain syndrome [16].

In experimental animals, Lewis, Wistar and Copenhagen rats have a high incidence of spontaneous nonbacterial prostatitis [14]. Administration of exogenous 17 β -estradiol can induce 100% of the incidence on prostatitis in old Wistar rats [18] and castration also has a

similar effect [13, 18]. Naslund et al. [13] reported that histopathological findings were very similar between spontaneous nonbacterial prostatitis and estradiol-induced prostatitis in rats [13]. These histopathological findings in rat spontaneous nonbacterial age-dependent prostatitis demonstrated several similarities to pathologically defined chronic prostatitis in human [19, 20]. These findings suggested that this rat model would be a useful model for the study of the treatment of human chronic prostatitis. Therefore, we decided to investigate the effects and mechanisms of CN-009 using a nonbacterial prostatitis rat model [13, 14] induced by 17 β -estradiol injection and castration.

No differences in the prostate weight were found in CN-009 630, CN-009 1260, T-60, and GBX




Fig. 7. Effects of CN-009 on stromal PCNA positive cell counts of the prostate. Each column represents the mean \pm SD.




Fig. 8. Effects on CN-009 on acinar epithelial TUNEL positive cell counts of prostate. Each column represents the mean \pm SD.

groups compared with the Control group. Since the weight of the prostate is mostly determined by the amount of residual secretory fluid, these findings may indicate that CN-009 cannot prevent the reduction of secretory prostatic fluid.

In the CN-009 1260 group, we observed roundly shaped acinar glands, cuboidal acinar epithelial cells containing secretory materials with positive PAP staining and diminished cell infiltrations into the lumen compared with the Control group. The acinar epithelial score was significantly increased. CN-009 could protect acinar epithelial function and cell shape against nonbacterial inflammation. GBX had a similar effect to CN-009 in the acinar glands. T-60 was not effective in the acinar epithelial function of this rat model. Therefore, GBX may play a central role for the

protection of epithelial damage in the NBP. The effect of GBX was similar to that of TS. However, GBX does not contain androgenic activity, because GBX has no effect on normal and hypertrophic prostate (unpublished data, 1968). Accordingly, this protective effect of GBX is discriminated from TS effect. In an in vitro study, GBX was reported to inhibit the cyclooxygenases and 5-lipoxygenases in the biosynthesis of the prostaglandins and leucotrienes [21]. Since prostaglandins and leucotriens enhance inflammatory cell infiltrations, GBX may protect against inflammation into the acinar lumen by inhibition of these enzymes. Furthermore, CN-009 showed an inhibition on the heat-induced hemolysis, which is correlated to lysosomal membrane stability [11]. CN-009 appears to




Fig. 9. Effects of CN-009 on stromal TUNEL positive cell counts of the prostate. Each column represents the mean \pm SD. **Significantly different from the Control group at $P < 0.01$.




Fig. 10. Effects of CN-009 on acinar epithelial score of the prostate. Each column represents the mean \pm SD. ##Significantly different from the Sham-ope group at $P < 0.01$. **Significantly different from the Control group at $P < 0.01$.

stabilize a lysosomal membrane, recover cell function and protect against degeneration of the acinar epithelium.

In addition, T-60 was shown to inhibit the growth of an immortal prostate cancer cell line in vitro [22]. However, their mechanisms are unknown. In the present study, the ratio of stromal area was significantly decreased in the CN-009 1260 and T-60 groups. Stromal TUNEL positive cell counts were increased in these groups. Therefore, CN-009 and T-60 may inhibit stromal cell proliferations by enhanced apoptosis. Although the exact mechanism of this process is unclear, several speculations are possible such as the direct effect by the apoptosis of fibroblast, and the indirect effect by the apoptosis of lymphocytes through the inhibition of several cytokines, such as several interleukins.

Further laboratory studies are necessary to elucidate the exact mechanisms of this compound.

Since no toxicological effects have been shown even in long-term administration, CN-009 is thought to be a safe drug [6, 23]. Here we reported the effects and mechanisms of CN-009 on rat experimental nonbacterial prostatitis model. CN-009 will also be a safe and effective against human nonbacterial chronic prostatitis.

In conclusion, CN-009 can work as a potent anti-inflammatory agent against chronic prostatitis. The present findings suggest that GBX, a fat soluble fraction of CN-009, protects the function and shape of acinar glandular epithelium and T-60, a water soluble fraction of CN-009, inhibits stromal cell proliferations in association with enhanced apoptosis.




Fig. 11. Effects of CN-009 on stromal ratio in the prostate. Each column represents the mean \pm SD. ##Significantly different from the Sham-ope. Group at $P < 0.01$. **Significantly different from the Control group at $P < 0.01$. *Significantly different from the Control group at $P < 0.05$.

Acknowledgements

We thank M. Komukai, M. Ishii, F. Kimura and E. Higaki for their technical assistance.

References

1. Brunner H, Weinder W, Scheifer HG. Studies on the role of *Ureaplasma urealyticum* and *Mycoplasma hominis* in prostatitis. *J Infect Dis* 1983;147:807-813.
2. Domingue GJ, Hellstrom WJ. Prostatitis. *Clin Microbiol Rev* 1999;11:2-203
3. Arakawa S, Matsui T, Gohji K, Okada H, Kamidono S. Prostatitis – the Japanese view point. *Int J Antimicrob Agents* 1997;11:623-646.
4. Donovan DA, Nicholas PK. Prostatitis; diagnosis and treatment in primary care. *Nurse Practitioner* 1997;22:144-156.
5. Pewitt EB, Schaeffer AJ. Urinary tract infection in urology; including acute and chronic prostatitis. *Infect Dis Clin North Am* 1997;11:623-646.
6. Jodai A, Maruta N, Shimomae E, Sakuragi T, Shindo K, Saito Y. A long term therapeutic experience with Cernilton in chronic prostatitis. *Acta Urol Jpn (Jpn.)* 1988;34:561-568.
7. Ito R, Ishii M, Yamashita S, Noguchi K, Kunikane K, Ohkubo Y, Tsushima Y, Sato S, Akamatsu H. Antiprosthetic hypertrophic action of Cernitin pollen-extract (Cernilton). *Pharmacometrics (Jpn.)* 1986;31:1-11.
8. Shah PJR. The treatment of disorders of the prostate with the rye-grass extract Prostabrit. *Comp Therp Med* 1996;4:21-25.
9. Onodera S, Yoshinaga M, Takenaga K, Toyoshima A, Uchiyama T. Effects of Cernitin pollen extract (CN-009) on the isolated bladder smooth muscles and the intravesical pressure. *Foliopharmacol Japan (Jpn.)* 1991;97:267-276.
10. Nakase K, Takenaga K, Hamanaka T, Kimura M. Inhibitory effect and synergism of Cernitin pollen extract on the urethral smooth muscle and diaphragm of the rat. *Folia pharmacol Japan (Jpn.)* 1988;91:385-392.
11. Ito R, Noguchi K, Yamashita S, Namikata S, Takenaga K, Yosinaga M, Shimuzu K, Ishii M, Mori N. Anti-inflammatory effects of Cernitin pollen extract (Cernilton). *Pharmacometrics (Jpn.)* 1984;28:55-65.
12. Ask-Upmark E. On a new treatment of prostatitis. *Grana Palynologica* 1960;2:115-118.
13. Naslund MJ, Strandberg JD, Coffey DS. The role of androgens and estrogens in the pathogenesis of experimental nonbacterial prostatitis. *J Urol* 1988;140:1049-1053.
14. Robinette CL. Sex-hormone-induced inflammation and fibromuscular proliferation in the rat lateral prostate. *The Prostate* 1988;12:271-286.
15. Schmidt JD, Patterson MC. Needle biopsy study of chronic prostatitis. *J Urol* 1966;96:519-533.
16. True LD, Berger RE, Rothman I, Ross SO, Krieger JN. Prostate histopathology and the chronic prostatitis/ chronic pelvic pain syndrome: a prospective study. *J Urol* 1999;162:2014-2018.
17. Aumuller G, Enderle-Schmitt U, Seitz J, Muntzig J, Chandler JA. Ultrastructure and immunohistochemistry of the lateral prostate on aged rats. *The Prostate* 1987;10:245-256.
18. Seethalakshmi L, Bala RS, Malhotra RK, Austin-riche T, Miller-Graziano C, Menon M, Lubernarod J. 17 B-estradiol induced prostatitis in the rat is an autoimmune disease. *J Urol* 1996;156:1838-1842.
19. Lundgren R, Holmquist B, Hesselvik M, Muntzig J. Treatment of prostatitis in the rat. *The Prostate* 1984;5:277-284.
20. Muntzig J, Sufrin G, Murphy GP. Prostatitis in the rat. *Scand J Urol Nephrol* 1979;13:17-22.
21. Loschen G, Ebeling L. Hemmung der Arachidonsaure-Kaskade durch einen Extrakt aus Roggenpollen. *Arzneim-Forsch Drug. Res* 1991;41:162-167.
22. Habib FK, Ross M, Buch AC, Ebeling L, Lewenstein: in vitro evaluation of the pollen extract, Cernitin T-60, in the regulation of prostate cell growth. *Br J Urol* 1990;66:393-397.
23. Rugendorff EW, Weinder W, Ebeling L, Buck AC. Results of treatment with Pollen extract (Cernilton N) in chronic prostatitis and prostatodynia. *Br J Urol* 1993;71:433-438.